DEPARTMENT OF HEALTH AND HUMAN SERVICES

NATIONAL INSTITUTES OF HEALTH

NATIONAL INSTITUTE OF ENVIRONMENTAL HEALTH SCIENCES

Hearing on³ ([D P L Q L Q J W K H) H G H U D O G R Z Q W R Z Q W K H U L V
S R O \ I O X R U R D O N \ O V X E V W D Q F H V 3) $ 6 ´

Testimony before the
Senate Committee on Environment and Public Works

Linda S Birnbaum, Ph.D., D.A.B.T., A.T.S.
Director, National Institute of Environmental Health Sciences and National Toxicology Program0 G [(Dire)7(

March 28, 2019

supported research uses human observational studies, animal models, *in vitro* tissue and cell culture systems, *in silico* approaches, and high throughput screening to study the effects of environmental exposure.

The most conclusive research focuses on a single chemical to understand the cause and effect on human health. While studying potentially toxic chemicals, we are largely limited to natural history and population-based studies that attempt to find connections between populations exposed and health effects in the real world. For that reason, you will hear me talk about [illegible] effects happened to more people than normal in populations that are exposed.

The research conducted to date reveals associations between PFAS exposures and a variety of specific adverse human health outcomes. These include the potential for [illegible] cognitive and neurobehavioral development, immune system dysfunction, endocrine disruption, obesity, diabetes, lipid metabolism, and cancer. While knowledge about these epidemiologic associations has steadily expanded in recent years, many questions remain unanswered. The NIEHS and NTP, in coordination with other Federal agencies and State and local governments, continue to conduct research to enhance our understanding of the potential mechanisms and biological processes through which PFAS may affect human health. NIEHS coordinates and participates in governmental health research to assure applicability, disseminate findings, and prevent duplication of effort. To this end, NIEHS continues to co-host and participate in numerous symposia and collaborative working groups.

Per and Polyfluoroalkyl Substances (PFAS)

Before detailing the health effects associated with PFAS exposures, it is necessary to describe this class of chemicals. First created in the 1930s and 1940s, PFAS include some 4,700 man-made chemicals that contain fluorine atoms bonded to a carbon chain.[4] The carbon-fluorine bond is one of the strongest ever created by man and is rarely seen in nature. That unique chemical composition of PFAS imparts desirable physical and chemical properties for consumer and industrial products, such as oil and water repellency, high and low temperature stability, and friction reduction. These properties have led to PFAS incorporation in a wide range of consumer products, including textiles, paper products, semiconductors, automotive and aerospace components, cookware, food packaging, and stain repellants. In addition, PFAS play an important role in

humans.

Decreased Immune System Function

As early as 1978, scientists observed immunotoxicity in non-human primates exposed to PFAS.[15] In 2016, NTP conducted a systematic literature review which included that PFOA and PFOS are presumed to be a hazard to healthy immune system function in humans.[16] This conclusion is based on a high level of evidence that PFOA and PFOS suppress antibody response in animal studies, and a moderate level of evidence that these chemicals affect multiple aspects of the immune system in humans. Adult PFAS exposure has also been associated with decreases in antibody production.[17] NTP is building on this 2016 systematic review to evaluate immunotoxicity of six related PFAS: PFDA, PFNA, PFHxA, PFBA, PFBS and PFHxS.[18]

Cancer

The epidemiological data on associations between PFAS and cancer risk are limited. Three published studies were recently summarized by the Agency for Toxic Substances and Disease Registry (ATSDR) in their Draft Toxicological Profile for Perfluoroalkyls.[19] According to the 7R[LFRORJLFDO 3URILOH [20] 2FFXSDWLRQDO DQG FRPPXQLW\ the risk of testicular and kidney cancer associated with PFOA. One consistent epidemiologic evidence for other cancer types were found for PFOA.[20,21] For PFOS, one occupational exposure study reported an increase in bladder cancer,[22] but this was not supported by subsequent occupational studies. General population studies have not consistently reported increases in malignant tumors for PFOS. Epidemiologic studies examining other perfluoroalkyl compounds consisted of two case-control studies. No increases in breast cancer risk were observed for PFHxS or PFNA; an increased breast cancer risk was observed for PFOSA.[23]

---

[15] Goldenthal EI, Jessup DC, Geil RG, Mehring JS. Final report, ninety day subacute rhesus monkey toxicity study, International Research and Development Corporation, study no. 91037, November 10, 1978, U.S. EPA Administrative Record, AR226-0447.

[16] National Toxicology Program. Monograph on Immunotoxicity Associated with Exposures to PFOA and PFOS. Sept. 2016. Research Triangle Park, NC: U.S. Internet: https://ntp.niehs.nih.gov/pubhealth/hat/noms/pfoa/index.html

[17] Kielsen K, Shamim Z, Ryder LP, Nielsen F, Grandjean P, Budtz-Jørgensen E, Heilmann C. Antibody Response to Booster Vaccination with Tetanus and Diphtheria in Adults Exposed to Perfluorinated Alkylates. *J. Immunotoxicol.* 2016; 13(2):276-83. DOI: 10.3109/1547691X.2015.106717

studies are consistent with and provide additional biological plausibility for the developmental effects observed in the human studies.[34,35]

Endocrine Disruption

Studies suggest that some PFAS may interfere with healthy hormonal function in the body. Our endocrine system controls our basic physiology, including how we metabolism, growth, fertility, and development. Human studies suggest a concern that early-life exposures to some PFAS may contribute to altered insulin resistance.[36,37] Although further confirmation is required, the findings from one study suggest that exposure to some PFAS during pregnancy may influence lipid metabolism and glucose tolerance.[38] A study of pregnant women in Cincinnati found that those with higher prenatal PFAS levels had children with higher body fat levels at age eight, a finding reinforced by other epidemiological studies[39,40,41] and similar effects on excessive body weight gain reported for experimental animals.[42] It appears that some PFAS may also affect body weight later in life. Scientists at the Harvard School of Public Health have found that adults with higher blood levels of some PFAS have lower resting metabolic rates, meaning they burn fewer calories while resting, which makes it difficult for them to maintain weight loss.[43]

---

[34] Valvi D, Oulhote Y, Weihe P, Dalgård C, Bjerve KS, Steuerwald U, Grandjean P. Gestational diabetes and offspring birth size at elevated environmental pollutant exposures.

awarded 10 new research project grants, representing a more than 30% increase in its

Leadership Summit hosted by EPA in May 2018.[86] Within the Department of Health and Human Services and primarily through NTP, NIEHS works closely with the FDA and CDC on PFAS matters. Additionally, NIEHS is specifically being consulted by ATSDR on the design and conduct of the exposure assessments and health studies authorized by the National Defense Authorization Act for Fiscal Year 2018, as amended.[87]

Conclusion

Thank you again for allowing me to share a scientific perspective on this important topic. In closing, I note that NIEHS is well positioned to continue contributing essential scientific knowledge about this complex and large class of chemicals. This knowledge can help regulators make sound, science-based decisions and informs the medical and public health communities about the potential health effects associated with exposure to PFAS. I welcome your questions.

https://factor.niehs.nih.gov/2018/3/science-highlights/pfas/index.htm.

[86] U.S. Environmental Protection Agency. EPA PFAS National Leadership Summit and Engagement. May 22, 2018. Internet: https://www.epa.gov/pfas/pfas-national-leadership-summit-and-engagement.

[87] Sec. 316 of the National Defense Authorization Act for Fiscal Year 2018, Public Law 115-91. December 12, 2017.

Bassler J, A Ducatman, M Elliott, S Wen, B Wahlang, J Barnett and MC Cave
Environmental perfluoroalkyl acid exposures are associated with liver disease characterized by apoptosis and altered serum adipocytokines [Journal Article]
Environ. Pollut. (2019) v. 247 pp. 1055-1063
Full-Text at: http://dx.doi.org/10.1016/j.envpol.2019.01.064

Brusseau ML
The influence of molecular structure on the adsorption of PFAS to fluid-fluid interfaces: Using QSPR to predict interfacial adsorption coefficients [Journal Article]
Water Research (2019) v. 152 pp. 148-158
Full-Text at: http://dx.doi.org/10.1016/j.watres.2018.12.057

Brusseau ML, N Yan, S Van Glubt, Y Wang, W Chen, Y Lyu, D Bingan, KC Carroll and FO Holguin
Comprehensive retention model for PFAS transport in subsurface systems [Journal Article]
Water Research (2019) v. 148 pp. 41-50
Full-Text at: http://dx.doi.org/10.1016/j.watres.2018.10.035

Cardenas A, R Hauser, DR Gold, KP Kleinman, MF Hivert, AF Fleisch, PD Lin, AM Calafat,

Harris MH, E Oken, SL Rifas-Shiman, AM Calafat, X Ye, DC Bellinger, TF Webster, RF White and SK Sagiv
Prenatal and childhood exposure to per-

Trevino LS and TA Katz
Endocrine Disruptors and Developmental Origins of Nonalcoholic Fatty Liver Disease
[Review]
Endocrinology (2018) v. 159 (1): pp. 201
Full-Text at: